

$N = 4$ Supersymmetric Quantum Mechanics with Magnetic Monopole

Soon-Tae Hong*

Department of Science Education, Ewha Womans University, Seoul 120-750 Korea

Joochan Lee†

Department of Physics, University of Seoul, Seoul 130-743 Korea

Tae Hoon Lee‡

Department of Physics, Soongsil University, Seoul 156-743 Korea

Phillial Oh§

*Department of Physics and Institute of Basic Science,
Sungkyunkwan University, Suwon 440-746 Korea*

(Dated: February 1, 2008)

We propose an $N = 4$ supersymmetric quantum mechanics of a charged particle on a sphere in the background of Dirac magnetic monopole and study the system in the $CP(1)$ model approach. By using the Dirac quantization method, we explicitly calculate the symmetry algebra taking the operator ordering ambiguity into consideration. We find that it is given by the superalgebra $su(1|2) \times su(2)_{\text{rot}}$. We also show that the Hamiltonian can be written in terms of the Casimir invariant of $su(2)_{\text{rot}}$ algebra. Using this relation and analyzing the lower bound for angular momentum, we find the energy spectrum. We, then, examine the ground energy sector to find that the $N = 4$ supersymmetry is spontaneously broken to $N = 2$ for certain values of the monopole charge.

PACS numbers: 11.30.Pb, 11.30.Qc, 14.80.Hv

Keywords: $N = 4$ supersymmetric quantum mechanics; magnetic monopole; $su(1|2) \times su(2)_{\text{rot}}$ algebra; supersymmetry breaking

Since the pioneering work of Dirac the quantum mechanics of a charged particle in the background of a Dirac magnetic monopole [1, 2] has attracted a great deal of attention due to its rich physical and mathematical properties. Its supersymmetric extension has been also proposed [3] and its various aspects has been studied [4, 5, 6, 7]. In the supersymmetric version, it was found [4] that the system originally written in $N = 1$ formulation [3] possesses an additional (hidden) supersymmetry, making the system in fact $N = 2$ supersymmetric. In Ref. [4] it was also pointed out that the hidden supercharge is related to the theory restricted to S^2 , the subspace of R^3 representing a fixed distance from the location of the monopole. On the other hand, manifest $N = 2$ superspace formulation of the system is possible on S^2 due to its Kähler structure, and the system confined to S^2 has been investigated in detail recently [6, 7]. In particular, in Ref. [6] the complete energy spectrum with the corresponding wave functions were found, and in Ref. [7] the issue of spontaneous symmetry breaking (whether or not the ground state is invariant under the supersymmetries) was discussed using the $CP(1)$ model approach. In this context, one of the interesting questions would be to ask whether the supersymmetry of the system could be extended further.

In this paper, we propose an explicit model of $N = 4$ supersymmetric quantum mechanics of a charged particle on a sphere in the background of Dirac magnetic monopole. Following the previous work on $N = 2$ supersymmetric quantum mechanics [7], we adopt the $CP(1)$ model approach. This allows $N = 4$ formulation of the system including the monopole interaction in component formalism. The dynamical variables consist of bosonic variables $(\bar{z}_i, z_j)$ and fermionic variables $(\bar{\psi}_{\alpha i}, \psi_{\beta j})$ ($i, j = 1, 2$; $\alpha, \beta = 1, 2$) satisfying $\bar{z} \cdot z - 1 = 0$, $\bar{z} \cdot \psi_{\alpha} = 0$, $\bar{\psi}_{\alpha} \cdot z = 0$. Note that the fermionic variables are doubled in number compared to the $N = 2$ case, as indicated by the internal indices, α, β . The bosonic variables are related to the space coordinate by the Hopf map $\vec{x} = \bar{z}\vec{\sigma}z$.

In quantizing the system there appears a parameter associated with the choice of operator ordering in defining the basic commutation relations as in the $N = 2$ case [7]. We study how physical quantities such as energy and angular momentum depend on this parameter. After quantization, we find that the symmetry algebra of our system is given by $su(1|2) \times su(2)_{\text{rot}}$. The bosonic $su(2)$ sector of $su(1|2)$ is the internal rotations associated with α, β indices, whereas the $su(2)_{\text{rot}}$ corresponds the angular momentum. Using the relations Hamiltonian with the supercharges and the the

*Electronic address: soonhong@ewha.ac.kr

†Electronic address: joochan@kerr.uos.ac.kr

‡Electronic address: thlee@ssu.ac.kr

§Electronic address: plohn@newton.skku.ac.kr

Casimir invariant of the rotational algebra we find the energy spectrum. We also investigate the ground state sector and find that the spontaneous supersymmetry breaking occurs for some particular values of the magnetic charge.

In order to proceed, we consider $N = 4$ supersymmetric transformations of the following form

$$\begin{aligned}\delta_\alpha z &= \psi_\alpha, & \delta_\alpha \bar{z} &= 0, & \delta_\alpha \psi_\beta &= 0, & \delta_\alpha \bar{\psi}_\beta &= 2i\nabla_{\alpha\beta}\bar{z}, \\ \bar{\delta}_\alpha z &= 0, & \bar{\delta}_\alpha \bar{z} &= \bar{\psi}_\alpha, & \bar{\delta}_\alpha \psi_\beta &= 2i\nabla_{\alpha\beta}z, & \bar{\delta}_\alpha \bar{\psi}_\beta &= 0,\end{aligned}\tag{1}$$

where

$$\nabla_{\alpha\beta}\bar{z} = \delta_{\alpha\beta}D_t\bar{z} - \frac{i}{2}(\bar{\psi}_\beta \cdot \psi_\alpha - \delta_{\alpha\beta}\bar{\psi}_\gamma \cdot \psi_\gamma)\bar{z},\tag{2}$$

and its complex conjugate

$$\nabla_{\alpha\beta}z = \delta_{\alpha\beta}D_t z + \frac{i}{2}(\bar{\psi}_\alpha \cdot \psi_\beta - \delta_{\alpha\beta}\bar{\psi}_\gamma \cdot \psi_\gamma)z.\tag{3}$$

The covariant derivative D_t is defined by

$$\begin{aligned}D_t z &= (\partial_t - ia)z, \\ D_t \bar{z} &= (\partial_t + ia)\bar{z},\end{aligned}\tag{4}$$

with a given by

$$a = -\frac{i}{2}(\bar{z} \cdot \dot{z} - \dot{\bar{z}} \cdot z) - \frac{1}{2}\bar{\psi}_\alpha \cdot \psi_\alpha.\tag{5}$$

Note a is invariant under the supersymmetric transformations of Eq. (1). One can check that the above supersymmetry transformations (1) preserve the supersymmetric $CP(1)$ constraints [8]

$$\bar{z} \cdot z - 1 = 0, \quad \bar{z} \cdot \psi_\alpha = 0, \quad \bar{\psi}_\alpha \cdot z = 0.\tag{6}$$

Our supersymmetric Lagrangian is then proposed by

$$L = 2|D_t z|^2 + \frac{i}{2}(\bar{\psi}_\alpha \cdot D_t \psi_\alpha - D_t \bar{\psi}_\alpha \cdot \psi_\alpha) - \frac{1}{4}((\epsilon_{\alpha\beta}\bar{\psi}_\alpha \cdot \psi_\beta)^2 + (\bar{\psi}_\alpha \cdot \psi_\alpha)^2) - 2ga,\tag{7}$$

which can be expressed in the following form;

$$\begin{aligned}L &= 2|\dot{z} - (\bar{z} \cdot \dot{z})z|^2 + \frac{i}{2}(\bar{\psi}_\alpha \cdot \dot{\psi}_\alpha - \dot{\bar{\psi}}_\alpha \cdot \psi_\alpha) - \frac{i}{2}(\bar{z} \cdot \dot{z} - \dot{\bar{z}} \cdot z)\bar{\psi}_\alpha \cdot \psi_\alpha \\ &\quad - \frac{1}{4}((\epsilon_{\alpha\beta}\bar{\psi}_\alpha \cdot \psi_\beta)^2 + (\bar{\psi}_\alpha \cdot \psi_\alpha)^2) + ig(\bar{z} \cdot \dot{z} - \dot{\bar{z}} \cdot z - i\bar{\psi}_\alpha \cdot \psi_\alpha).\end{aligned}\tag{8}$$

In the above Lagrangian, we put the electric charge $e = -1$, and g is the magnetic monopole charge. Compared to the previous $N = 2$ case [7], it has additional quartic fermionic interaction terms which are essential for the existence of $N = 4$ supersymmetry.

Next, we perform the canonical quantization of the system. We define the momenta p and $\bar{p}$ conjugate, respectively, to the fields z and $\bar{z}$,

$$p = 2D_t\bar{z} + \frac{i}{2}(\bar{\psi}_\alpha \cdot \psi_\alpha + 2g)\bar{z}, \quad \bar{p} = 2D_t z - \frac{i}{2}(\bar{\psi}_\alpha \cdot \psi_\alpha + 2g)z.\tag{9}$$

The classical Hamiltonian is

$$H_c = 2(D_t\bar{z})(D_t z) - g\bar{\psi}_\alpha \cdot \psi_\alpha + \frac{1}{4}((\epsilon_{\alpha\beta}\bar{\psi}_\alpha \cdot \psi_\beta)^2 - (\bar{\psi}_\alpha \cdot \psi_\alpha)^2).\tag{10}$$

It should be supplemented by the following six second class constraints

$$C_1 = \bar{z} \cdot z - 1, \quad C_2 = p \cdot z + \bar{z} \cdot \bar{p}, \quad C_{3\alpha} = \bar{z} \cdot \psi_\alpha, \quad C_{4\alpha} = \bar{\psi}_\alpha \cdot z,\tag{11}$$

and one first class constraint,

$$C_0 = -i(\bar{z} \cdot \bar{p} - p \cdot z) - \bar{\psi}_\alpha \cdot \psi_\alpha + 2g,\tag{12}$$

the Gauss law constraint corresponding to the local $U(1)$ symmetry. Classically, two quartic terms in the fermion field in Eq. (10) cancel each other. We keep it here because they produce a quadratic term when quantized. We quantize this theory following the Dirac scheme. We start with the Poisson bracket relations

$$\{z_i, p_j\} = \{\bar{z}_i, \bar{p}_j\} = \delta_{ij}, \quad \{\bar{\psi}_{i\alpha}, \psi_{j\beta}\} = -i\delta_{ij}\delta_{\alpha\beta}, \quad (13)$$

with the remaining brackets being zero. To incorporate the second class constraints, we calculate the Dirac brackets using the definition given by

$$\{A, B\}_D = \{A, B\} - \{A, C_a\}\Theta^{ab}\{C_b, B\}, \quad (14)$$

where Θ^{ab} is the inverse matrix of $\Theta_{ab} = \{C_a, C_b\}$. From this result we obtain the following quantum commutation (and anti-commutation) relations upon replacing $\{A, B\}_D \rightarrow -i[A, B]$,

$$\begin{aligned} [p_i, z_j] &= -i\delta_{ij} + \frac{i}{2}\bar{z}_i z_j, & [p_i, \bar{z}_j] &= \frac{i}{2}\bar{z}_i \bar{z}_j, \\ [p_i, p_j] &= \frac{i}{2}(p_i \bar{z}_j - p_j \bar{z}_i), & [\bar{p}_i, p_j] &= \frac{i}{2}(\bar{z}_j \bar{p}_i - z_i p_j) - \alpha\psi_{i\alpha}\bar{\psi}_{j\alpha} + \beta\bar{\psi}_{j\alpha}\psi_{i\alpha}, \\ [\bar{\psi}_{i\alpha}, \psi_{j\beta}] &= \delta_{\alpha\beta}(\delta_{ij} - \bar{z}_i z_j), & [p_i, \bar{\psi}_{j\alpha}] &= i\bar{\psi}_{i\alpha}\bar{z}_j, \end{aligned} \quad (15)$$

with $\alpha + \beta = 1$. The above brackets are supplemented by their Hermitian conjugates, and the remaining commutators all vanish. They form a straightforward generalization of $N = 2$ case [7]. In the second line, the operator ordering in the second bracket is chosen by the condition that the dynamical variables commute with the second class constraint, C_2 , ordered as $p \cdot z + \bar{z} \cdot \bar{p} = 0$. Note that this does not fix the operator ordering completely and we still have undetermined α (or β) in Eq. (15) as in the $N = 2$ case [7].

In order to obtain the symmetry algebra of the system, we compute the Noether charges associated with various global symmetries. The space rotations are generated by

$$\begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \rightarrow e^{-\frac{i}{2}w^a\sigma^a} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}, \quad (16)$$

whose operator-ordered conserved charge is given by

$$K_a = \frac{i}{2}\bar{z}\sigma_a\bar{p} - \frac{i}{2}p\sigma_a z + \gamma\bar{z}\sigma_a z + \frac{1}{2}\bar{\psi}_\alpha\sigma_a\psi_\alpha. \quad (17)$$

Here we have added the third term associated with the operator ordering ambiguity. We find that K_a 's generate the required rotation and satisfy the $SU(2)$ algebra

$$[K_a, K_b] = i\epsilon_{abc}K_c, \quad (18)$$

provided the following conditions are satisfied

$$\alpha = \frac{1}{4} + \gamma, \quad \beta = \frac{3}{4} - \gamma. \quad (19)$$

The Noether charge associated with the phase symmetry of the fermionic variables yields the conserved charge

$$N_F = \bar{\psi}_\alpha \cdot \psi_\alpha. \quad (20)$$

The supercharges are given by

$$Q_\alpha = p \cdot \psi_\alpha, \quad \bar{Q}_\beta = \bar{\psi}_\beta \cdot \bar{p}. \quad (21)$$

Note that the supercharges have no ordering ambiguity. The internal $SU(2)$ rotations define isospin operators S^a by

$$S^a = \frac{1}{2}\bar{\psi}_\alpha\sigma^a_{\alpha\beta}\psi_\beta. \quad (22)$$

A straightforward calculation yields the following relations

$$S^2 = -\frac{3}{4}N_F^2 + \frac{3}{2}N_F = -\frac{3}{4}(N_F - 1)^2 + \frac{3}{4}, \quad (23)$$

where $S^2 = S^a S^a$. Denoting the eigenvalue of S^2 by $s(s+1)$, we find the relations

$$\begin{aligned} s = 0 & \leftrightarrow N_F = 0, \\ s = \frac{1}{2} & \leftrightarrow N_F = 1, \\ s = 0 & \leftrightarrow N_F = 2, \end{aligned} \quad (24)$$

For local $U(1)$ symmetry we choose as the generator

$$G_0 \equiv -i(\bar{z} \cdot \bar{p} - p \cdot z) - \bar{\psi}_\alpha \cdot \psi_\alpha. \quad (25)$$

Classically, this quantity being equal to minus the twice of the monopole charge is the Gauss law constraints, Eq. (12). To use this constraint to select the physical Hilbert space, however, one should take care of the ordering problem. This amounts to adding an appropriate constant before setting G_0 equal to $-2g$. As our quantum mechanical Hamiltonian we choose the following expression

$$\begin{aligned} H &= \frac{1}{2}(p \cdot \bar{p} - p \cdot z \bar{z} \cdot \bar{p}) - \frac{i}{2}(\bar{z} \cdot \bar{p} - p \cdot z) \bar{\psi}_\alpha \cdot \psi_\alpha - (\alpha + \frac{1}{4}) \bar{\psi}_\alpha \cdot \psi_\alpha \\ &= \frac{1}{2}(p \cdot \bar{p} - p \cdot z \bar{z} \cdot \bar{p}) + \frac{1}{2} \left(G_0 - 2\alpha + \frac{1}{2} \right) \bar{\psi}_\alpha \cdot \psi_\alpha + \frac{1}{2} (\bar{\psi}_\alpha \cdot \psi_\alpha)^2 - \frac{1}{2} \bar{\psi}_\alpha \cdot \psi_\alpha. \end{aligned} \quad (26)$$

One can show that the above Hamiltonian commutes with the supercharges. It can be obtained by quantizing the classical Hamiltonian of Eq. (10) if $G_0 - 2\alpha + 1/2$ is identified with $-2g$. This strongly suggests that the aforementioned ordering constant is $-2\alpha + \frac{1}{2}$ and $G_0 - 2\alpha + \frac{1}{2}$ should be interpreted as the monopole charge. However, for consistency of this interpretation we must show that $G_0 - 2\alpha + \frac{1}{2}$ is quantized according to the Dirac quantization condition of the magnetic monopole charge. It will be shown shortly (see Eq. (39) and discussions below) that this is indeed the case. Still there is an ambiguity of adding an integer in the choice of the above mentioned ordering constant. A different choice of this integer will correspond to a different theory (or a different interpretation of the theory). In this paper we write the quantum mechanical Gauss law constraint as

$$G_0 - 2\alpha + \frac{1}{2} = -2\tilde{g}, \quad (27)$$

with $\tilde{g}$ being treated as the monopole charge and our physical states are required to satisfy the above Gauss law constraint.

A complete algebraic structure of the symmetry generators can be computed to yield

$$\begin{aligned} [S^a, S^b] &= i\epsilon_{abc} S^c, & [S^a, N_F] &= 0, \\ [Q_\alpha, N_F] &= Q_\alpha, & [\bar{Q}_\alpha, N_F] &= -\bar{Q}_\alpha, \\ [S^a, Q_\alpha] &= -\frac{1}{2} \sigma_{\alpha\beta}^a Q_\beta, & [S^a, \bar{Q}_\alpha] &= \frac{1}{2} \bar{Q}_\beta \sigma_{\beta\alpha}^a, \\ [Q_\alpha, Q_\beta] &= 0, & [\bar{Q}_\alpha, \bar{Q}_\beta] &= 0, \\ [Q_\alpha, \bar{Q}_\beta] &= 2H\delta_{\alpha\beta} - 2\tilde{g} \left(S^a \sigma_{\alpha\beta}^a - \frac{1}{2} N_F \delta_{\alpha\beta} \right). \end{aligned} \quad (28)$$

Also the K_a operators in Eq. (17) satisfy,

$$\begin{aligned} [K_a, K_b] &= i\epsilon_{abc} K_c, & [K_a, N_F] &= 0, \\ [K_a, Q_\alpha] &= 0, & [K_a, \bar{Q}_\alpha] &= 0. \end{aligned} \quad (29)$$

Defining $\tilde{N}_F \equiv N_F + 2H/\tilde{g}$ and $q_\alpha \equiv Q_\alpha/\sqrt{-2\tilde{g}}$, $\bar{q}_\beta \equiv \bar{Q}_\beta/\sqrt{-2\tilde{g}}$ ($\tilde{g} < 0$) we find the last equation of Eq. (28) becomes

$$[q_\alpha, \bar{q}_\beta] = -\frac{1}{2} \tilde{N}_F \delta_{\alpha\beta} + S^a \sigma_{\alpha\beta}^a. \quad (30)$$

Other commutation relations remain the same because $[H, S^a] = [H, N_F] = [H, Q_\alpha] = [H, \bar{Q}_\beta] = 0$ and we find that $K_a, S^a, \tilde{N}_F, q_\alpha$ and $\bar{q}_\beta$ generate $su(1|2) \times su(2)_{\text{rot}}$ symmetry [9]. (The case of $\tilde{g} > 0$ can be covered by redefining $S^a \rightarrow \frac{1}{2} \bar{\psi}_\alpha (-\sigma_{\alpha\beta}^{aT}) \psi_\beta$, $N_F \rightarrow -\bar{\psi}_\alpha \cdot \psi_\alpha$ and interchanging $Q_\alpha \rightarrow \bar{Q}_\alpha$, $\bar{Q}_\alpha \rightarrow Q_\alpha$.)

In order to examine the energy spectrum let us write the Hamiltonian in terms of the supercharges, as

$$H = \frac{1}{4}[Q_\alpha, \bar{Q}_\alpha]_+ - \frac{1}{2}\tilde{g}N_F, \quad (31)$$

where we have added the subscript $+$ to emphasize that the bracket is the anti-commutator. This relation can be obtained by taking the trace of the last line of Eq. (28). Since the first term is non-negative one finds that the energy is bounded from below by the second term in the above equation. Note that each sector with a definite fermion number has a different bound. However, whether or not these bounds can be saturated depends on the existence of the states in each sector that are invariant under the supersymmetry. In other words they may not be the optimal bounds. In order to obtain the true energy bound for each sector for a given parameter $\tilde{g}$, we proceed as follows. First, note that

$$\begin{aligned} p \cdot \bar{p} - p \cdot z \bar{z} \cdot \bar{p} &= p_i(\delta_{ij} - z_i \bar{z}_j) \bar{p}_j \\ &= p_i \epsilon_{ij} \bar{z}_j \ z_l \epsilon_{kl} \bar{p}_k \equiv a \bar{a} \end{aligned} \quad (32)$$

where we have defined

$$a = \epsilon_{ij} p_i \bar{z}_j, \quad \bar{a} = \epsilon_{kl} z_l \bar{p}_k. \quad (33)$$

They satisfy the following commutation relation

$$[a, \bar{a}] = -G_0 + 2\alpha + \frac{3}{2} - 2\bar{\psi}_\alpha \cdot \psi_\alpha = 2(\tilde{g} - \Sigma), \quad (34)$$

where $\Sigma \equiv N_F - 1$ can be regarded as the total spin along the radial direction, and Eq. (27) was used. Eigenvalues of the spin operator, Σ , consists of three values, $\sigma = +1, 0, -1$, reflecting that we have two spin half degrees of freedom. Next, we need the bound for $a\bar{a}$. If $\tilde{g} - \sigma \leq 0$, the bound is zero, and the state saturating this bound is obtained by imposing $\bar{a} = 0$. If $\tilde{g} - \sigma \geq 0$, $a\bar{a}$ is bounded by $2(\tilde{g} - \sigma)$ as one can see from $a\bar{a} = \bar{a}a + 2(\tilde{g} - \sigma)$ and $\bar{a}a \geq 0$. In the latter case the bound is saturated by the states satisfying $a = 0$. These two cases can be combined into

$$a\bar{a} \geq |\tilde{g} - \sigma| + (\tilde{g} - \sigma). \quad (35)$$

This result together with Eqs. (26), (27) yields the bound for the energy

$$E_{\min} = \frac{1}{2}(|\tilde{g} - \sigma| + (\tilde{g} - \sigma)) - \tilde{g}(\sigma + 1) + \frac{1}{2}(\sigma + 1)^2 - \frac{1}{2}(\sigma + 1) \quad (36)$$

$$= \frac{1}{2}|\tilde{g} - \sigma|(|\tilde{g} - \sigma| + 1) - \frac{1}{2}\tilde{g}(\tilde{g} + 1). \quad (37)$$

By comparing this minimum energy with $-\frac{1}{2}\tilde{g}N_F$ of Eq. (31) we conclude that the following types of supersymmetric ground states exist: $\sigma = -1$ if $\tilde{g} \leq -1$, $\sigma = 0$ if $\tilde{g} = 0$, and $\sigma = 1$ if $\tilde{g} \geq 1$.

One can obtain further information from the relation between the Hamiltonian and the angular momentum squared

$$H = \frac{1}{2}(K^2 - \tilde{g}(\tilde{g} + 1)). \quad (38)$$

Comparing this with Eq. (37) leads to the minimum value of the angular momentum quantum number


$$k_{\min} = |\tilde{g} - \sigma|. \quad (39)$$

Since the angular momentum quantum number must be half-integer or integer and the spin σ is an integer, $\tilde{g}$ must also be a half-integer or an integer. In particular, one finds that $k_{\min}$ is a half-integer if $\tilde{g}$ is a half-integer, and an integer if $\tilde{g}$ is an integer. This confirms that the quantization of $\tilde{g}$ and $\tilde{g}$ can indeed be interpreted as the effective monopole charge. More generally, the angular momentum quantum number can be written as

$$k = k_{\min} + n, \quad (n = 0, 1, 2, \dots) \quad (40)$$

and the energy spectrum can be written as

$$E = \frac{1}{2}k(k + 1) - \frac{1}{2}\tilde{g}(\tilde{g} + 1). \quad (41)$$

FIG. 1: Diagram for k versus $\tilde{g}$.

We give the diagram for k versus $\tilde{g}$ in Fig. 1. Among the states satisfying Eq. (39), the ground states are represented by the points marked with dots and circles. The dots correspond to the supersymmetric ground states for given values of $\tilde{g}$. The cases, $\tilde{g} = \pm 1/2$, are somewhat special. In these cases there is no $N = 4$ supersymmetric states. In other words, there is no state killed by both Q_α and $\bar{Q}_\alpha$. For $\tilde{g} = 1/2$, for instance, the ground states, marked with a circle in Fig. 1, consist of two sectors, $(k = 1/2, \sigma = 0)$ and $(k = 1/2, \sigma = 1)$, each of which has a twofold angular momentum degeneracy. However, one can argue in this case that the ground state still has $N = 2$ supersymmetry left over. Note that the supercharges commute with the Hamiltonian and the angular momentum. Thus, applying the supercharges to any state will not change the energy and angular momentum quantum numbers. On the other hand the commutation relations $[\Sigma, Q_\alpha] = -Q_\alpha$ and $[\Sigma, \bar{Q}_\alpha] = \bar{Q}_\alpha$ tell us that $Q_\alpha, \bar{Q}_\alpha$ play the role of lowering and raising operators of the fermion number. Now, let $|\psi\rangle$ be a ground state with $\sigma = 0$. Then $Q_\alpha|\psi\rangle = 0$ because there is no $\sigma = -1$ state to be mapped in the ground energy sector. Similarly, $\bar{Q}_\alpha|\psi\rangle$ belongs to the $\sigma = 1$ sector of the lowest energy level. Simple counting together with the fact that the supercharges commute with K_a suggests that a certain linear combination of $\bar{Q}_\alpha|\psi\rangle$ should vanish. To conclude, for $\tilde{g} = \pm 1/2$ the ground state is invariant under the half of the supersymmetry.

In summary, we have shown that the quantum mechanics of a charged particle on a sphere in the background of Dirac magnetic monopole allows $N = 4$ supersymmetric extension. Using the Dirac quantization procedure, we explicitly calculated the symmetry algebra and found that it is given by the superalgebra $su(1|2) \times su(2)_{\text{rot}}$. We also investigate the spectrum of the Hamiltonian which is given by the Casimir invariant of the $su(2)_{\text{rot}}$ symmetry. By analyzing the ground energy sector we found that the supersymmetry is spontaneously broken for particular values of $\tilde{g} = \pm \frac{1}{2}$. There are a few unresolved aspects deserving a further study: The first one is to look for a manifest superspace formulation of our system by using the $N = 4$ chiral superfield. The other is the construction of the complete wavefunctions by explicitly realizing Eq. (15) as differential operators on an appropriate function space. Some of these are currently under progress.

Acknowledgments

STH would like to acknowledge financial support in part from the Korea Science and Engineering Foundation grant (R01-2000-00015). PO was supported by Korea Research Foundation grant (R05-2004-000-10682-0). The authors

would like to thank the ATCTP for the hospitality during their visit.

-
- [1] P. A. M. Dirac, *Proc. Roy. Soc. Lon.* **A 133** (1931) 60.
 - [2] S. R. Coleman, *The Magnetic Monopole Fifty Years Later* in Les Houches Sum. School (1981) 461; P. Goddard and D. I. Olive, *Rep. Prog. Phys.* **41** (1978) 1357; R. Jackiw, *Ann. Phys.* **129** (1980) 183.
 - [3] E. D'Hoker and L. Vinet, *Phys. Lett.* **B 137** (1984) 72; *Lett. Math. Phys.* **8** (1984) 439.
 - [4] F. De Jonghe, A. J. Macfarlane, K. Peeters, and J. W. van Holten, *Phys. Lett.* **B 359** (1995) 114, hep-th/9507046.
 - [5] G. W. Gibbons, R. H. Rietdijk and J. W. van Holten, *Nucl. Phys.* **B 404** (1993) 42, hep-th/9303112; D. Spector, *Phys. Lett.* **B 474** (2000) 331, hep-th/0001008; G. Papadopoulos, *Class. Quantum Grav.* **17** (2000) 3715, hep-th/0002007; M. S. Plyushchay, *Phys. Lett.* **B 485** (2000) 187, hep-th/0005122; C. Leiva and M. S. Plyushchay, *Phys. Lett.* **B 582** (2004) 135, hep-th/0311150.
 - [6] S. Kim and C. Lee, *Ann. Phys.* **296** (2002) 390, hep-th/0112120 and references therein.
 - [7] S. T. Hong, J. Lee, T. H. Lee and P. Oh, *Phys. Rev.* **D 72** (2005) 015002, hep-th/0505018.
 - [8] A. D'Adda, P. Di Vecchia and M. Lüscher, *Nucl. Phys.* **B 152** (1979) 125 and references therein.
 - [9] A. Pais and V. Rittenberg, *J. Math. Phys.* **16** (1975) 2062; P. G. O. Freund and I. Kaplansky, *J. Math. Phys.* **17** (1976) 228; M. Scheunert, W. Nahm and V. Rittenberg, *J. Math. Phys.* **18** (1977) 155; M. Marcu, *J. Math. Phys.* **21** (1980) 1277.